arXiv:1605.04677v1 [math.AC] 16 May 2016

AN EASY PROOF OF THE GENERAL NERON DESINGULARIZATION IN DIMENSION ≤ 1

ASMA KHALID AND ZUNAIRA KOSAR

Abstract. In this paper we give an easy proof of the General Neron Desingularization in the frame of regular morphisms between Artinian local rings and Noetherian local rings of dimension one.
Key words : Flatness, Completion, Smooth morphisms, regular morphisms.
2010 Mathematics Subject Classification: Primary 13B40, Secondary 14B25,13H05,13J15.

Introduction

A ring morphism $u : A \to A'$ has *geometrically regular fibers* if for all prime ideals $P \in Spec\ A$ and all finite field extensions K of the fraction field of A/P the ring $K \otimes_{A/P} A'/PA'$ is regular. A flat morphism u is *regular* if its fibers are geometrically regular. A regular morphism of finite type is *smooth* and a localization of a smooth algebra is *essentially smooth*.

Let $(A, \mathfrak{m})$ be a Noetherian local ring of dimension ≤ 1, A' a Noetherian local ring of dimension one and $u : A \to A'$ a regular morphism. Let $B = A[Y]/I$, $Y = (Y_1, \ldots, Y_n)$. Then any morphism $v : B \to A'$ factors through a smooth A-algebra C, that is v is a composite A-morphism $B \to C \to A'$. This is a particular case of the General Neron Desingularization given in [4], [8], [5], with a very difficult proof. An algorithmic proof is given in [3] for the case when A, A' are domains, and it is extended in [7] for arbitrary Noetherian local rings of dimension one.

Let $(A, \mathfrak{m})$ be a local Artinian ring, $(A', \mathfrak{m}')$ a Noetherian complete local ring of dimension one such that $k = A/\mathfrak{m} \cong A'/\mathfrak{m}'$, and $u : A \to A'$ be a regular morphism. Suppose that $k \subset A$. Then $\bar{A}' = A'/\mathfrak{m}A'$ is a discrete valuation ring (shortly a DVR). Choose $x \in A'$ such that its class modulo $\mathfrak{m}A'$ is a local parameter of $\bar{A}'$, that is, it generates $\mathfrak{m}'\bar{A}'$. Let $B = A[Y]/I$, $Y = (Y_1, \ldots, Y_n)$.

The purpose of this paper is to give an easy proof to the following theorem.

Theorem 1. *Then any morphism $v : B \to A'$ factors through a smooth A-algebra C, that is, v is a composite A-morphism $B \to C \to A'$.*

The proof is illustrated in Example 7.

1. Preliminaries

A ring morphism $h : A \to B$ is called *quasi-smooth* [8] if for any A-algebra D and an ideal $I \subset D$ with $I^2 = 0$, any A-morphism $B \to D/I$ lifts to an A-morphism

We gratefully acknowledge the support from the ASSMS GC. University Lahore, for arranging our visit to Bucharest, Romania and we are also greatful to the Simion Stoilow Institute of the Mathematics of the Romanian Academy for inviting us.

$B \to D$. A finitely presented quasi-smooth morphism is smooth. Let $u : A \to B$ be a quasi smooth morphism and C an A-algebra. Then $C \otimes u$ is also quasi smooth which is known as the *Base change* property.

Let A be a DVR, x a local parameter of A, $A' = \hat{A}$ its completion and $B = A[Y]/I$, $Y = (Y_1, \ldots, Y_n)$ a finite type A-algebra. If $f = (f_1, \ldots, f_r)$, $r \leq n$ is a system of polynomials from I then we consider an $r \times r$-minor M of the Jacobian matrix $(\partial f_i/\partial Y_j)$. Let $c \in N$. Suppose that there exist an A- morphism $v : B \to A'/(x^{2c+1})$ and $N \in ((f) : I)$ such that $v(NM) \notin (x)^c/(x^{2c+1})$, where for simplicity we write $v(NM)$ instead $v(NM + I)$.

Theorem 2. [6] *There exists a B-algebra C which is smooth over A such that every A-morphism $v' : B \to A'$ with $v' \equiv v$ modulo x^{2c+1} (that is $v'(Y) \equiv v(Y)$ modulo x^{2c+1}) factors through C.*

Corollary 3. *Any A-morphism $v : B \to A'$ factors through a smooth A-algebra C.*

This corollary follows also from the classical Néron desingularization [2].

Theorem 4. [7] *Let $u : A \to A'$ be a regular morphism of one dimensional Noetherian local rings and B a finite type A-algebra. Suppose that A contains its residue field. Then any A-morphism $v : B \to A'$ factors through a smooth A-algebra C, that is v is a composite A-morphism $B \to C \to A'$.*

2. Proof of Theorem 1 when $A = k$

Then A' is a complete DVR and has the form $A' = k[[x]]$. Let $A_1 = k[x]_{(x)}$ and u_1 the inclusion $A_1 \subset A'$. Then u_1 is flat because A_1 is principal and A' has no A-torsion. The fraction field extension induced by u_1 is separable and A' is the completion of A_1. It follows that u_1 is regular because A_1 is excellent (see e.g. [1]). Let be as above, B a k-algebra of finite type, and $v : B \to A'$ an A-morphism. Let $B_1 = A_1 \otimes_A B$ and $v_1 : B_1 \to A'$ be the composite map $a_1 \otimes b \mapsto u_1(a_1) \cdot v(b)$.

Lemma 5. *Suppose that v_1 factors through a smooth A_1-algebra $\tilde{C}$ then v factors through a smooth A-algebra C.*

Proof. Since $B_1 = A_1 \otimes_k B = A_1[Y]/IA_1[Y]$ where $Y = (Y_1, \ldots, Y_n)$ and v_1 can be factored through a smooth A_1-algebra $\tilde{C}$, we get the following the commutative diagram:



where $\delta : A_1 \to B_1$ is given by $a_1 \mapsto a_1 \otimes 1$, $a_1 \in A_1$, $\gamma : B \to B_1$ is given by $b \mapsto 1 \otimes b$, $b \in B$ and v_1 is the composite map $B_1 \xrightarrow{\beta} \tilde{C} \xrightarrow{w} A'$. Then $\tilde{C}$ has the form $(A_1[Z]/(f_1, \ldots, f_r))_{M\tilde{h}}$, $Z = (Z_1, \ldots, Z_s)$ for some $r \times r$-minor M of the

Jacobian matrix $\partial f/\partial Z$ and some $\tilde{h} \in A_1[Z]$ such that $w(\tilde{h})$ is invertible in A'. Clearly $\tilde{C}$ is essentially smooth over k but unfortunately not smooth because it is not a k-algebra of finite type. Now choose $h \in k[x] \setminus (x)$ such that $f_1, \ldots, f_r$ have coefficients in $k[x, Z]_h$. Set $E = (k[x, Z]_h/(f_1, \ldots, f_r))_{M,\tilde{h}}$, then $\tilde{C}$ is a localization of E with respect to all $g \in k[x] \setminus (x)$. Thus $\tilde{C}$ is the filtered inductive limit of E_g, $g \in k[x] \setminus (x)$. Let $\varphi_g : E_g \to \tilde{C}$ be the limit maps.

We claim that $\alpha = \beta\gamma$ factors through a certain E_g. Indeed, let $\alpha(\hat{Y}_i) = q_i/g_i$ for some $q_i \in E$ and $g_i \in k[x] \setminus (x)$. Then α factors through E_g for $g = \prod_{i=1}^n g_i$. Clearly v is the composite map $B \to C = E_g \xrightarrow{\varphi_g} \tilde{C} \xrightarrow{w} A'$, that is, v factors through the smooth k-algebra C. □

The following proposition is closed to [6, Corollary 16].

Proposition 6. *Suppose that $A = k$, then v factors through a smooth k-algebra C.*

Proof. By Corollary 3, v_1 factors through a smooth $A_1 = k[x]_{(x)}$-algebra $\tilde{C}$. Let's say v_1 is the composite map $B_1 \to \tilde{C} \xrightarrow{w} A'$. Then by Lemma 5, v factors through a smooth k-algebra C. □

3. The Proof of Theorem 1

Since A is an Artinian local ring there exists a certain s such that $\mathfrak{m}^s = 0$. A has the form $A = k[T]/\mathfrak{a}$, $T = (T_1, \ldots, T_m)$, and the maximal ideal of A is generated by T. Then for all $i \in [m] = \{1, \ldots, m\}$, ${T_i}^s \in \mathfrak{a}$ and $A' = k[[x]][T]/(a) \cong A \otimes_k k[[x]]$. Suppose that $B = A[Y]/I$ where $Y = (Y_1, \ldots, Y_n)$, $v : B \to A'$ and $v(\hat{Y}_i) = \hat{y}_i = \sum_{\alpha \in \mathbb{N}^m, |\alpha| < s} y_{i\alpha} \cdot T^\alpha$, $T^\alpha = {T_1}^{\alpha_1} \cdots {T_m}^{\alpha_m}$, $|\alpha| = \alpha_1 + \cdots + \alpha_m \ < s$ and $y_{i\alpha} \in k[[x]]$. We have the following commutative diagram.

$$\begin{array}{ccccccc} k & \longrightarrow & \bar{B} & \longrightarrow & \bar{B}_1 & \xrightarrow{\bar{v}_1} & k[[x]] \\ \downarrow & & & & & & \uparrow \\ A & \longrightarrow & A \otimes_k \bar{B} & \longrightarrow & A \otimes_k \bar{B}_1 & \longrightarrow & A \otimes_k k[[x]] = A' \end{array}$$

where $\bar{B} = B/\mathfrak{m}B$. Set $\overline{B_1} = k[(y_{i\alpha})_\alpha] \subset k[[x]]$ and let $\overline{v_1}$ be this inclusion. Then v factors through $B_1 = A \otimes_k \bar{B}_1 \subset A'$, that is v is the composite map $B \xrightarrow{q} B_1 \xrightarrow{A \otimes_k \bar{v}_1} A'$, where q is defined by $Y_i \to \sum_\alpha T^\alpha \otimes y_{i\alpha}$. Applying the Proposition 6, we see that $\bar{v}_1$ factors through a smooth k-algebra $\bar{C}$. Then $A \otimes_k \bar{v}_1$ factors through $C = A \otimes_k \bar{C}$ and so v factors through C.

Example 7. Let $A = \mathbf{Q}[t]/(t^2)$, $A' = \mathbf{Q}[[x]][t]/(t^2)$, $A_1 = \mathbf{Q}[x]_{(x)}[t]/(t^2)$, $B = A[Y_1, Y_2]/(Y_1^3 - Y_2^2)$ and u_1, v_1 two formal power series from $\mathbf{Q}[[x]]$ which are algebraically independent over $\mathbf{Q}(x)$ and $u_1(0) = v_1(0) = 1$. By the Implicit Function Theorem there exists $u_2 \in k[[x]]$ such that $u_2^2 = u_1^3$. Set $v_2 = (3/2)xu_1^2v_1u_2^{-1}$, $\hat{y}_1 = x^2u_1 + tv_1$, $\hat{y}_2 = x^3u_2 + txv_2$. We have $f(\hat{y}_1, \hat{y}_2) = x^6(u_1^3 - u_2^2) + tx^4(3u_1^2v_1 - 2u_2v_2) = 0$ and we may define $v : B \to A'$ by $Y \to (\hat{y}_1, \hat{y}_2)$. Take $\overline{B_1} = \mathbf{Q}[x, x^2u_1, x^3u_2, v_1, xv_2]$. Then $\bar{v} = \mathbf{Q} \otimes_A v : B/tB \to \mathbf{Q}[[x]]$ factors through $\overline{B_1}$. Let $\varphi : \mathbf{Q}[x, Y_{10}, Y_{20}, Y_{11}, Y_{21}] \to$

$\overline{B_1}$ be given by $Y_{10} \to x^2u_1$, $Y_{20} \to x^3u_2$, $Y_{11} \to v_1$, $Y_{21} \to xv_2$. Then $\operatorname{Ker}\varphi$ contains $g_1 = Y_{10}^3 - Y_{20}^2$, $g_2 = 3Y_{10}^2Y_{11} - 2Y_{20}Y_{21}$. The Jacobian matrix $\partial g_k/\partial Y_{ij}$ contains a 2×2-minor $M = 4Y_{20}^2$ which is mapped by φ in $4x^6u_2^2$. Taking $c = 7$ we see that $\varphi(M) \not\equiv 0$ modulo x^c. Note that φ induces the isomorphism $F = \mathbf{Q}[x, Y_{10}, Y_{20}, Y_{11}, Y_{21}]/(g_1, g_2) \cong \overline{B_1}$ because $\operatorname{Ker}\varphi = (g_1, g_2)$ since $\dim F = \dim \overline{B_1} = 3$, u_1, v_1 being algebraically independent over $\mathbf{Q}(x)$.

Take $E = \mathbf{Q}[x, u_1, u_2, v_1, v_2]$. The kernel of the map $\psi : \mathbf{Q}[x, U_1, U_2, V_1, V_2] \to E$ contains the polynomials $h_1 = U_1^3 - U_2^2$, $h_2 = 3U_1V_1 - 2U_2V_2$ and as above we see that $\mathbf{Q}[x, U_1, U_2, V_1, V_2]/(h_1, h_2) \cong E$. Now the Jacobian matrix $\partial h/\partial(U_i, V_i)$ contains a minor $M' = 4U_2^2$ which is mapped by ψ in $4u_2^2 \notin (x)$. Then

$$\bar{C} = (\mathbf{Q}[x, U_1, U_2, V_1, V_2]/(h_1, h_2))_{V_2U_2U_1} \cong (\mathbf{Q}[x, U_1, V_1, V_2]/(U_1^3 - (9/4)U_1^2V_1^2V_2^{-2}))_{V_2U_1}$$
$$\cong (\mathbf{Q}[x, U_1, V_1, V_2]/(U_1 - (9/4)V_1^2V_2^{-2}))_{V_2} \cong \mathbf{Q}[x, V_1, V_2]_{V_2}$$

is smooth over $\mathbf{Q}$ and the inclusion $\bar{v}_1 : \overline{B_1} \to \mathbf{Q}[[x]]$ factors through $\bar{C}$ because $\bar{v}_1$ is the composite map

$$\overline{B_1} \cong \mathbf{Q}[x, Y_{10}, Y_{20}, Y_{11}, Y_{21}]/(g_1, g_2) \xrightarrow{\rho} \bar{C} \xrightarrow{\psi} E \to \mathbf{Q}[[x]],$$

where ρ is given by $Y_{10} \to x^2U_1$,$Y_{20} \to x^3U_2$, $V_i \to V_i$. Then $C = A \otimes_{\mathbf{Q}} \bar{C}$ is smooth over A and v factors through C because it is the composite map $B \xrightarrow{\nu} B_1 = A \otimes_{\mathbf{Q}} \overline{B_1} \xrightarrow{A\otimes_{\mathbf{Q}}\bar{v}_1} A \otimes_{\mathbf{Q}} \mathbf{Q}[[x]] \cong A'$, where ν is given by $Y_1 \to (1 \otimes x^2u_1 + t \otimes v_1)$, $Y_2 \to (1 \otimes x^3u_2 + t \otimes xv_2)$.

References

[1] H. Matsumura, *Commutative Algebra*, Benjamin, New-York, 1980.

[2] A. Néron, *Modeles minimaux des varietes abeliennes sur les corps locaux et globaux*, Publ. Math. IHES, **21**, 1964, 5-128.

[3] A. Popescu, D. Popescu, *A method to compute the General Neron Desingularization in the frame of one dimensional local domains*, to appear in "Singularities and Computer Algebra - Festschrift for Gert-Martin Greuel, On the Occasion of his 70th Birthday", Editors Wolfram Decker, Gerhard Pfister, Mathias Schulze, Springer Monograph., arXiv:AC/1508.05511.

[4] D. Popescu, *General Neron Desingularization and approximation*, Nagoya Math. J., **104** (1986), 85-115.

[5] D. Popescu, *Artin Approximation*, in "Handbook of Algebra", vol. 2, Ed. M. Hazewinkel, Elsevier, 2000, 321-355.

[6] D. Popescu, *Around General Neron Desingularization*, to appear in Journal of Algebra and Its Applications, **16**, No. 2 (2017), arXiv:1504.06938.

[7] G. Pfister, D. Popescu, *Constructive General Neron Desingularization for one dimensional local rings*, arXiv:AC/1512.08435v1.

[8] R. Swan, *Neron-Popescu desingularization*, in "Algebra and Geometry", Ed. M. Kang, International Press, Cambridge, (1998), 135-192.

Abdus Salam School of Mathematical Sciences,GC University, Lahore, Pakistan.
E-mail address: asmakhalid768@gmail.com
E-mail address: zunairakosar@gmail.com